


## Theory Question 2: Solution:

### Nuclear Masses and Stability

(a) The alpha-decay process is as follows:



Therefore the energy criterion for decay to happen is:

$$m_A - m_{A-4} - m_4 > 0$$

The number and type of nucleons in the decay is preserved so we only have to consider the binding energies:

$$-B_A + B_{A-4} + B_4 > 0$$

If we write  $B/A = a + bA$ , where  $a$  and  $b$  are constants to be found from the graph, then this equation becomes:

$$-A(a+bA) + (A-4)(a+b(A-4)) + B_4 > 0$$

$$-8bA - 4a + 16b + B_4 > 0$$

By inspecting the graph, a good linear approximation to  $B/A$  above  $A = 100$  is:

$$B/A = (9.6 - 0.0080 \times A) \text{ MeV}$$

i.e.  $a = 9.6$  MeV and  $b = 0.0080$  MeV, and the condition becomes:

$$-0.064A - 38.4 - 0.1 + 25.0 > 0$$

$$\underline{A > 13.5/0.064 = 211}$$

Part (b).

(i) Because  $A$  is fixed we only need to consider the penultimate two terms which depend on  $Z$ .

$$\frac{dB}{dZ} = -2Za_c A^{-1/3} - \frac{a_a}{A} (-4A + 8Z)$$

$$Z_{\max} = \frac{4a_a}{2a_c A^{-1/3} + 8a_a / A} = \frac{A}{2} \frac{1}{\left(1 + \frac{a_c A^{2/3}}{4a_a}\right)}$$

(ii)  $Z_{\max} = 79.25$

The full expression for the differential equation in (a) is:

$$\frac{dB}{dZ} = -2Za_c A^{-1/3} - \frac{a_a}{A}(-4A + 8Z) \pm 2a_p A^{-3/4}$$

The last term is positive if a change in  $Z$  of +1 changes the nucleus from an even-even to an odd-odd, and negative if the reverse is true. Note  $A$  is positive in this case.

How do we deal with the last term?

The number  $Z_{\max}$  has to be an integer, and even numbers are favoured over odd so we can guess  $Z_{\max} = 80$ . To check, evaluate the last three terms for various values of  $Z$ :

77	979.241
78	975.915
79	976.295
80	975.341
81	978.093
82	979.512
83	984.637

This confirms that  $Z_{\max} = 80$ ; this is an even-even nucleus.


(iii) Consider only the last three terms in the equation; the rest are constant if  $A$  is constant. Call the sum of these quantities  $X$ . To find out whether these nuclei are stable we need to find differences in  $X$  between neighbouring species and to compare these differences with the energy requirements for each would-be decay process.

(i)  $\beta^-$  - decay;  $n \rightarrow p + e^-$ , need  $\Delta X > -1.30 + 0.51 = -0.79$  MeV

(ii)  $\beta^+$  - decay;  $p \rightarrow n + e^+$ , need  $\Delta X > 1.30 + 0.51 = 1.81$  MeV

(iii)  $\beta^-\beta^-$  - decay;  $2n \rightarrow 2p + 2e^-$ , need  $\Delta X > 2(-1.30 + 0.51) = -1.58$  MeV

(iv) Electron capture;  $e^- + p \rightarrow n$ , need  $\Delta X > 1.30 - 0.51 = 0.79$  MeV



↑ For information: graph not expected from students

Nucleus/Process	$X$ (MeV)
$^{128}_{51}\text{Sb}$	496.59
$^{128}_{52}\text{Te}$	491.19
$^{128}_{53}\text{I}$	491.06
$^{128}_{54}\text{Xe}$	489.16
$^{128}_{55}\text{Cs}$	492.54

↑ For information: table not expected from students

Nucleus/Process	$\beta^-$ - decay	$\beta^+$ - decay	Electron-capture	$\beta^-\beta^-$ - decay
$^{128}_{53}\text{I}$	√	0	0	√
$^{128}_{54}\text{Xe}$	0	0	0	0
$^{128}_{55}\text{Cs}$	0	√	√	0

Students will fill out this table

**Theory Question No.2: Mark Distribution**

Smallest fractional mark allowed: 0.25

Marks allowed for errors consistently propagated only if physically reasonable.

(a)	Approach	1.5	3
	Correct Answer	1.5	
(b)(i)	Approach	1	2
	Correct Answer	1	
(b)(ii)	Approach	1	2
	Correct Answer	1	
(b)(iii)	0.25 for each of 12 entries	3	3
	Grand Total		10